

## PRODUCTION REQUIREMENTS FOR 1 HECTARE OF MAIZE

Seed requirements: Generally require about 25 kgs of seed, specifically varies with variety, planting time, farming region

Plant population: 45 000 to 55 000 depending with the variety, time of planting, agro-ecological region, intended purpose eg fodder

Fertilizer requirements		Product	Rate/ha	Nitrogen	Phosphate	Potash	Sulphur	Comments	Rate/ha
Base Dressing	Agricura Compound D	7:14:7 9S	400	28	56	28	36	All at or before planting	
Top dressing	Ammonium Nitrate (34.5% N)	AN	400	138	0	0	0	Split apply at 3-4 & 6-7	
	OR		Total	166	56	28	36	weeks after planting	
Base dressing	Agricura Double D	14:28:14 1.8S	200	28	56	28	4	As above	
Top dressing	Ammonium nitrate (34.5% N)	AN	400	138	0	0	0	As above	
			Total	166	56	28	4		

## Herbicides, Insecticides and Fungicides

Problem	Product	Applicati <mark>on ra</mark> te App	plication Remarks			
Grasses,	S- <mark>Met</mark> olachlor/Dual Mag <mark>num</mark>	1.3 - 1.9 lt/ha	Pre-emergent against grasses and some broad leaved weeds			
B/L weeds,	A <mark>tran</mark> ex/Atrazine	3.6-5.5 lt/ha	Pre-emergent or early post against broad leaved weeds			
Sedges,	R <mark>elay/</mark> Harness/Acetochlo <mark>r</mark>	1 lt/ha	Pre-emergent against grasses and some broad leaves			
	Basagran/Bentazone	3.0- 5.0 lt/ha	Post-emergent against broad leaves and nut sedge			
	Halosulfuron	50 g/ha	Post-emergent. Grasses, B/L weeds & Yellow nutsedge.			
	Stellar Star	0.7-1.0 lt/ha	Post-emergent. Annaual grasses & some B/Lweeds.			
AI	MCPA	3 - 5 lt/ha	Pre or post emergent against broad leaved weeds only			
	Dicamba/Banvel	300 - 500mls/ha	Post emergent against broad leaved weeds only			
	Alachlor/Laso	3 - 4lt/ha	Pre-emergent against grasses and some B/L weeds			
	Nicosulfron/Ascent	45grms/ha	Post emergent against Shamva grass, grasses and B/Leaves			
	Terbutryn/terbutrex	2 - 4lt/ha	Pre-emergent against Grasses and B/L weeds			
	Paraquat	1 - 2lt/ha	Post emergent directed from tasseling			
Conservation						
Tillage	Roundup/Glyphosate	1 - 6 lt/ha	Apply to weeds before or just after planting			



Stalk borers	Lambda cyalothrin	200mls/ha	Apply as at first sign of infestation				
African Armyworm	Decis forte	125mls/ha	Apply as at first sign of infestation				
Cutworms	Carbaryl 85 WP	1kgs/ha	Full cover spray and repeat as necessary.				
Boll Worms	C <mark>hlop</mark> yrifos	350 - 400 mls	Spray over the top in 200 - 500 litres of water				
Fall Armyworm	Fenvelerate	200mls/ha	Apply as at first sign of infestation				
	Belt	38mls/ha	Apply as at first sign of infestation				
	C <mark>arta</mark> p hydrochloride	800grms/ha	Apply as at first sign of infestation				
	A <mark>vaun</mark> t/Steward	250mls/ha	Apply as at first sign of infestation				
	Karate zeon	200mls/ha	Apply as at first sign of infestation				
Dipterex/Kombat  Acephate/Orthene		3 - 4kgs/ha	Apply at the funnel of the plant within the first 3-5 weeks  Apply as at first sign of infestation				
		650grms/ha					
	Lannate/Methomyl/Methomex	300grms/ha	Apply as at first sign of infestation				
Hoppers /	Dimethoate 40 EC	500-750mls/ha	Apply as a high volume spray. Repeat every 7-10 days.				
Streak Virus	Gaucho 600FS	580mls/100kgs of seed	Apply as seed dresser at planting				
Grey leaf	Propiconazole 25EC	750mls/ha	Spray when disease is seen on 3rd leaf below the cob.				
spot	Tebuconazole	1lt/ha	Apply when disese first appears on basal 4 - 5 leaves				
	Benomyl/Benlate	500-750grms/ha	Leaves above the cob should be kept disease free.				
Termites,	Chlopyrifos	1lt/ha	Band apply at planting or spray 8 - 10 weeks after planting				
white grubs	Imidacloprid	200 mls	Apply 8 - 10 weeks after planting				
Stored grain	Actellic gold dust	25g/50kg bag (shelled grain)	Mix well into grain				
Insects							

Consult Agricura Agronomy team for technical information on herbicide combinations (tank mixtures) and insecticide combinations.

Approximately 12.5 x 16 litre knapsacks will cover 1 hectare and apply 200 litres of water Approximately 14 x 15 litre knapsacks will cover 1 hectare and apply 200 litres of water

ALWAYS READ LABELS BEFORE USING THE PRODUCTS

Head office landline: 04-621570-9, website: www.agricura.co.zw, email: agricura@agricura.co.zw, Agricura facebook page name: Agricura PVT Limited